AD-A237 286

1	
11	ו ר
\ <	/-

SECURITY CLASSIFICATION OF THIS PAGE					
REPORT	N rage	1619 HOUR IINI 15616 HSI	1 0///0	oved Ivu. v704-0188	
la REPORT SECURITY CLASSIFICATION	ECTE MAN	16. RESTRICTIVE N	MARKINGS		
20 SECURITY CLASSIFICATION AUTHORITY JUN	12 4 1991 7		AVAILABILITY OF	REPORT	
26. DECLASSIFICATION/DOWNGRADING SCHEDU	LE	Unlim	ited		
4. PERFORMING ORGANIZATION REPORT NUMBER	R	5. MONITORING C	RGANIZATION REI	PORT NUMBER(S)	
NA .		NA			
Washington University ST. Louis	6b. OFFICE SYMBOL (If applicable) NA	t	Javal Rese	earch	
6c. ADDRESS (City, State, and ZIP Code) 8 0 8 115 Washington University School 4566 Sco77 Aur S7. Lon	lof Medicino is MO 63110	ARUNGT	State, and ZIPCO PUINUTS	22217-	טטס'
8a. NAME OF FUNDING/SPONSORING ORGANIZATION	8b. OFFICE SYMBOL (If applicable)	9 PROCUREMENT	INSTRUMENT IDE	NTIFICATION NU	MBER
ONR -	ONR	N 000 14	-8814	-0032	
8c. ADDRESS (City, State, and ZIP Code) C. f. 76		10. SOURCE OF FU	JNDING NUMBERS	TASK	WORK UNIT
(,4. +5	•	ELEMENT NO	NO.	NO 441m803	ACCESSION NO.
11 TITLE (Include Security Classification)		<u> </u>	<u> </u>		L
an classified			,		
12. PERSONAL AUTHOR(S)					
13a. TYPE OF REPORT 13b TIME CO	OVERED TO	14. DATE OF REPOF	RT (Year, Month, D	Day) 15 PAGE	COUNT
16 SUPPLEMENTARY NOTATION	-141				
17 COSATI CODES	18. SUBJECT TERMS (C	Continue on reverse	of necessary and	identify by block	number)
FIELD GROUP SUB-GROUP	Vestibular		micicula		
	efferent,				
19 ABSTRACT (Continue on reverse if necessary and identify by block number) Sec a Hickard					
20 DISTRIBUTION / AVAILABILITY OF ABSTRACT UNCLASSIFIED/UNLIMITED SAME AS R	PT	21. ABSTRACT SEC	URITY CLASSIFICA	TION	
22a NAME OF RESPONSIBLE INDIVIDUAL	S C C OTENS	22b TELEPHONE (In	nclude Area Code)	22c OFFICE SY	MBOL

ABSTRACT

All vertebrates are andowed with a vestibular efferent system (EVS) consisting of somata within the Central Nervous System with long axons exiting the brain to innervate the labyrinth. Behaviorally relevant stimuli related to feeding and/or aggressive behaviors and conditions leading to enhanced attentional states or alerting, activate the EVS. Increased EVS activity modifies the resting rate and response dynamics to motion of vestibular afferents. This modification is non-uniform across the fiber spectrum of the semicircular canals, for example, affecting the more sensitive, low spontaneous activity cells more profoundly than their less sensitive counterparts. The cellular bases for EVS affects are excitatory axo-axonic synapses upon primary afferents and axo-somatic inhibitory synapses upon hair cells.

DTIG	COPY, INSPECTED

Acqua	sion For	
X11	TAASD	Ø
DIL	ran	
Unicidat	วายยุคส์	
Justi	fleation.	
By	· Carrier Committee Commit	
	trutton/	
Avai	bility	Cočes
	Avail and/or	
Dist	Special	
0 1		
H-1]	

FINAL REPORT ON CONTRACT NUMBER NOO014-0032

PRINCIPLE INVESTIGATOR Stephen M. Highstein

CONTRACT TITLE: Vestibular Efferent Activity in Squirrel Monkeys

START DATE 1, October, 1987 END DATE 1, October, 1990

RESEARCH OBJECTIVE:

To investigate the structure and function of the mammalian efferent vestibular system in Squirrel monkeys in order to aid in the understanding of the purpose for, and the potential utilization of this system. Intracellular recording will be taken from efferent vestibular neurons in anesthetized animals. We will stimulate the VIIIth nerves bilaterally to antidromically identify these neurons. Subsequently, we will identify sources of synaptic input to these neurons by electrically stimulating sites in the central and peripheral nervous systems and recording the post-synaptic potentials thereby evoked. We will inject horseradish peroxidase into these cells and into the efferent nucleus to determine relevant morphological features of the system. We will record extracellularly in alert animals from the somata of antidromically identified efferent vestibular neurons to define the level of spontaneous activity, if any, and to determine the responses of these neurons to behaviorally applied stimuli. Immunohistochemistry will also be employed to determine which peptide transmitterr candidates are present within the system.

PROGRESS (Years 1-3)

We have purchased the Cambridge Electronic Design recording system and the Asyst language to program this interface. We have written the programs to use this equipment in the proposed experiments and have begun the alert animal series proposed. We have recorded from three monkeys and are continuing to this line of investigation. Part of these experiments included the training of a post-doctoral fellow and a student who were unfamiliar with this type of recording. We have also performed several intracellular experiments in anesthetized, paralyzed animals. To date we have collected six efferent vestibular neurons. These cells receive input from both VIIIth nerves and the spinal cord. These experiments were also performed with an M.D.-Ph.D. student and an Otolaryngology resident as a part of their training in my laboratory. We have published several papers and abstracts listed below.

FUBLICATIONS AND REPORTS (Year 3):

- An oral presentation has been given at an ONR workshop in Pensacola this last winter. Several publications bearing on the efferent control of the labyrinth are listed below.
- 1) Tricas, T., and Highstein, S.M. The action of the efferent vestibular system upon the lateral line afferents in free swimming toadfish. Neurosci. abs., 1989.
- 2) Tricas, T., and Highstein, S.M. The action of the efferent vestibular system upon the lateral line afferents in free swimming toadfish. Biological Bulletin abs., 1989.
- 3) Boyle, R. and Highstein, S. M. Resting discharge and response dynamics of horizontal semicircular canal afferents of the toadfish, Opsanus tau. J. Neurosci., 10(5): 1557-1569, 1990.
- 4) Boyle, R. and Highstein, S. M. Efferent vestibular system in the toadfish: action upon horizontal semicircular canal afferents. J. Neurosci., 10(5): 1570-1582, 1990.
- 5) Carey, J. P., and Highstein, S. M., Physiology and brainstem morphology of single horizontal semicircular canal afferents in the toadfish, Opsanus tau: A biotin dye study. Neurosci. abs., 1990, in press.
- 5) Locke, R. E. and Highstein, S. M., Efferent modulation of synaptic noise in lagenar afferents of the toadfish. Neurosci. abs., 16:735, 1990.
- 7) Locke, R. E. and Highstein, S. M., Efferent modulation of synaptic noise and spike frequency in lagenar afferents of the toadfish. Association for Research in otolaryngology abs., 99, 1990.
- 8) Locke, R. E. and Highstein, S. M., Efferent modulation of synaptic noise and spike frequency in lagenar afferents of the toadfish. Biological Bulletin abs., 1989.
- 9) Tricas, T. and Highstein, S.M., Visual activation of the octavolateralis efferent system during predation in free swimming tradfish, Opsanus tau. Exp. Brain Res., 83:233-236, 1990.
- 10) Tricas, T. and Highstein, S.M. Action of the octavolateralis afferent system upon the lateral line of free-swimming toadfish, Opsanus 1964. J. Comp. Physiol., in press.
- 30yle, R., Carey, J.P., and Highstein, S.M. Morphological correlates of response dynamics and efferent stimulation in horizontal semicircular taxal afferents of the toadfish, Opsanus tau. J. Neurophysiol. in press.
- (2) Schassel, D.A., Ginzberg, R., and Highstein, G.M. Morphophysiology of synaptic transmission between type I hair nells and vestibular

primary afferents. An intracellular study employing horseradish peroxidase in the lizard, <u>Calotes versicolor</u>.

TRAINING ACTIVITIES: I have trained an Otolaryngology fellow in aspects of the techniques used in the published studies.